

The 1 μm discontinuity in the Extragalactic Background Light spectrum: an artefact of foreground subtraction

K. Mattila^{*}

Observatory, Tähtitorninmäki, P.O. Box 14, FI-00014 University of Helsinki, Finland

Accepted ; Received ; in original form

ABSTRACT

Several recent papers claim the detection of a near infrared Extragalactic Background Light (EBL) intensity at 1.25 - 4 μm that exceeds the integrated light of galaxies by factors of > 3 . When combined with a claimed optical detection of the EBL at 0.80 μm the EBL excess emission spectrum has a discontinuity at $\sim 1\mu\text{m}$. This discontinuity has given rise to an interpretation in terms of ultraviolet radiation emanating from the first generation of massive stars at redshifts of 7 - 20 (so called Population III stars). The interpretation of the NIR excess emission as being of extragalactic origin depends crucially on the model used in the subtraction of the Zodiacal Light, the dominant foreground contaminant. We estimate the Zodiacal Light at 0.80 μm using on the one hand the measurement by Bernstein, Freedman & Madore (2002b), with corrections for some omitted effects of atmospheric scattering and calibration, and on the other hand the model of Kelsall et al. (1998). There is in neither case any evidence for a step in the EBL at $\sim 1\mu\text{m}$. We emphasize that in order to avoid systematic effects it is essential to use the same Zodiacal Light model (Kelsall et al. 1998) for both the NIR (1.25 - 4 μm) and optical (0.80 μm) data. We emphasize, however, that our analysis does not allow a statement on the *overall level* of the NIR EBL. The contribution of the Diffuse Galactic Light to the “EBL excess” emission is estimated. It is found to be significant at 3 - 4 μm and should be carefully evaluated in future measurements which aim at detecting an EBL signal at the level of $\sim 10 \text{ nW m}^{-2} \text{ sr}^{-1}$, i.e. at the level of the integrated light of (known) galaxies.

Key words: cosmology: diffuse radiation - observations - infrared: general

1 INTRODUCTION

The Extragalactic Background Light (EBL) in the ultraviolet, optical, and infrared wavelength regions can potentially provide a measure for the total luminous matter in the Universe since the epoch of star and galaxy formation; for a review see Longair (2001).

Detection of a near infrared Extragalactic Background Light (EBL) component, in excess of the integrated light from known galaxies, has recently been announced by several research groups using the *COBE*/DIRBE (for a review see Dwek, Arendt, & Krennrich 2005) and the Japanese *IRTS* instruments (Matsumoto et al. 2005). A “first detection” of the optical EBL at 0.30, 0.55, and 0.80 μm has been claimed by Bernstein, Freedman & Madore (2002a,b) (hereafter referred to as BFM02a,b). Combining the claimed detections in these two wavelength bands, i.e. at 1.25 - 4 μm and 0.8 μm , there appears to be an indication of a discontinuity at $\sim 1\mu\text{m}$, the EBL brightness rising by a factor of

~ 4 from the short-wavelength to the long-wavelength side of this step (see e.g. Matsumoto et al. 2005). The potential importance of this spectral discontinuity stems from its suggested interpretation as signature of the first generation of stars (s.c. Population III stars) with their rest-frame emission at $\lambda > 0.912\mu\text{m}$ redshifted by a factor of $1+z \approx 8-20$ to the near infrared (Santos, Bromm, & Kamionkowski 2002; Salvaterra & Ferrara 2003; Cooray & Yoshida 2004; Madau & Silk 2005; Dwek, Arendt, & Krennrich 2005).

The reality of the NIR EBL excess at 1 - 4 μm and the 1 μm spectral step have been contested both on observational (see e.g. Aharonian et al. 2006; Dwek, Arendt, & Krennrich 2005) and on theoretical grounds (see e.g. Madau & Silk 2005). The present author (Mattila 2003) has argued that, because of insufficient or incorrect atmospheric scattering corrections and calibration, the claim by BFM02 for the detection of the EBL at 0.30, 0.55, and 0.80 μm was premature.

In the present paper we pose the question whether the BFM02 measurement can be used to set a sufficiently low *upper limit* to the EBL at 0.80 μm which, in combination

^{*} E-mail: mattila@cc.helsinki.fi

with the claimed NIR EBL values at 1.25 - 4 μm , would justify the claim for a step near $\sim 1\mu\text{m}$ in the EBL spectrum. Because of the large intensity of the Zodiacal Light in comparison with the EBL its accurate subtraction is crucial for any statement on the EBL and its spectral shape. Therefore, we examine in detail the underlying ZL determinations and models used in the above mentioned works. We will also examine the contribution of the Diffuse Galactic Light (DGL) to the observed 0.8 - 4 μm sky brightness.

2 ZODIACAL LIGHT ESTIMATION

The basic formula used in the above mentioned papers for extracting the NIR or optical EBL is the following:

$$I_{\text{EBL}} = I_{\text{tot}} - I_{\text{ZL}} - I_{\text{ISL}} - I_{\text{DGL}} \quad (1)$$

where I_{tot} is the total sky brightness as measured from space, I_{ZL} the Zodiacal Light (ZL) as estimated using dedicated observations or a model of the interplanetary dust cloud, I_{ISL} is the Integrated Starlight (ISL) from unresolved stars evaluated from star count models, and I_{DGL} is the Diffuse Galactic Light (DGL).

The first two terms are large, $\sim 300 \text{ nW m}^{-2} \text{ sr}^{-1}$ at 1 - 2 μm , as compared to the EBL which according to e.g. Matsumoto et al. (2005) and Cambr  sy et al. (2001) is $\sim 50\text{--}70 \text{ nW m}^{-2} \text{ sr}^{-1}$, but has conventionally been considered to be much smaller, $\sim 10 \text{ nW m}^{-2} \text{ sr}^{-1}$, and mainly due to the integrated light of galaxies only. While the ISL term is smaller, it is still several times larger than the EBL signal due to known galaxies. Thus, the demands for the accuracy of the I_{tot} and I_{ZL} determinations are extremely high.

2.1 ZL estimated from the observations of Bernstein, Freedman & Madore (2002a,b)

In the case of the claimed EBL detections at 0.30, 0.55, and 0.80 μm by BFM02 the I_{ZL} term was derived from ground based observations of the depths of Fraunhofer lines between 0.39 and 0.51 μm . Large uncertainties are caused to ground based night sky photometry by the corrections needed for the atmospheric scattered light and extinction. Also, the method necessitates to calibrate the *extended source* responses of two different telescopes, a ground based and a space borne one, to the same *absolute* scaling.

In a re-discussion of the BFM02 ZL observations Mattila (2003) showed that corrections had to be applied because of the following errors or omissions: (1) incorrect aerosol albedo; (2) omission of ground reflectance; (3) omission of DGL as source of atmospheric scattering; (4) incorrect aperture-correction factor. All these corrections were negative in sign and thus had the effect of decreasing the ZL value. Their combined effect to $I_{\text{ZL}}(0.80 \mu\text{m})$ amounted to -9.4 to $-13.1 \cdot 10^{-9} \text{ erg/s cm}^2 \text{ sr } \text{\AA}$, see Table 1 of Mattila (2003). This correction is ca. 10 times larger than the accuracy of $\pm 1 \cdot 10^{-9} \text{ erg/s cm}^2 \text{ sr}$ claimed by BFM02a for their ZL value and ca. 4-6 times as large as their claimed EBL value of $2.2 \cdot 10^{-9} \text{ erg/s cm}^2 \text{ sr } \text{\AA}$ at 0.80 μm . Each one of the corrections (1)–(4) has large uncertainties caused by the ill-defined atmospheric, ground reflectance and telescope PSF properties. We estimate that the resulting correction has an uncertainty of $\sim \pm 50\%$ or $\sim \pm 5 \cdot 10^{-9} \text{ erg/s cm}^2 \text{ sr } \text{\AA}$. In an

Erratum paper Bernstein, Freedman & Madore (2005) discussed a number of errors and omissions in BFM02. While some of the errors noted by Mattila (2003) were addressed, they did not yet address the four major corrections (1) - (4) as discussed above. The original ZL estimate of BFM02b is given in line 2 of Table 1 and the (Mattila 2003) estimate, including these corrections (1) - (4), in line 4 of Table 1.

Another point of concern is how BFM02 handled the *combination* of the individual systematic errors: this was done by assuming a flat probability distribution for each contributing source of error. This method produces combined systematic errors which are by factor 2 - 3 smaller than the conventionally determined errors obtained by adding in quadrature the individual systematic errors. One specific systematic error, important for the present purpose of estimating I_{ZL} at 0.80 μm , is caused by the reddening of the ZL relative to the Solar colour. BFM02a estimate the reddening correction factor per 1000 $\text{\AA}$ to be $C(\lambda) = 1.044$ with a systematic error of $[-0.014, +0.006]$. Extrapolating I_{ZL} from the reference wavelength of 0.465 μm to 0.80 μm the systematic error caused by the uncertainty in the ZL reddening correction thus becomes $[-4.1\%, +1.75\%]$. Combining this *in quadrature* with the other systematic errors the total systematic uncertainty of $I_{\text{ZL}}(0.8\mu\text{m})$ becomes $[-4.4\%, +2.4\%]$ or $[-3.0, +1.6] \cdot 10^{-9} \text{ erg/s cm}^2 \text{ sr } \text{\AA}$ (line 3 of Table 1).

2.2 ZL estimated using the Kelsall et al. (1998) model

The Zodiacal Light model which has been adopted in many of the NIR EBL studies, e.g. Matsumoto et al. (2005) and Cambr  sy et al. (2001), has been constructed by Kelsall et al. (1998). This model uses the full COBE/DIRBE data set at 10 wavelengths between 1.25 and 240 μm and relies essentially on the seasonal variations of the ZL. The IPD emission dominates the sky brightness at high galactic latitude areas and mid IR wavelengths, $\lambda \approx 10 - 60\mu\text{m}$, a situation which helps to constrain the model parameters at these wavelengths.

In order to investigate whether the different approaches used for the ZL estimation at $\lambda \approx 1\text{--}4\mu\text{m}$ (modelling) and at $\lambda = 0.80\mu\text{m}$ (ground based observations, BFM02b) could be the reason for the $\sim 1\mu\text{m}$ EBL spectrum step we have applied the Kelsall et al. (1998) ZL model to the I_{tot} observation of BFM02a at 0.80 μm . To do this we have extracted from the DIRBE Sky and Zodi Atlas (DSZA) the ZL value at J band (1.25 μm) for the specific sky position ($\lambda - \lambda_{\odot}$, β) and day of the year of the HST measurement of I_{tot} by BFM02a. This ZL value, in units of MJysr^{-1} , is given in line 2 of Table 2. The systematic error estimate given in brackets corresponds to the error of $15 \text{ nW/m}^2 \text{ sr}^{-1}$ as given in Table 7 of Kelsall et al. (1998). To transform the tabulated DSZA value into a monochromatic intensity at 1.25 μm a small colour correction is applied assuming for the ZL a solar temperature blackbody spectrum.

To estimate the ZL intensity at 0.80 μm we adopt the Solar SED as given by Colina, Bohlin, R.C.& Castelli (1996), convolve it with the HST WFPC2 filter F814W transmission curve using the SYNPHOT/calcpht procedure¹, and

¹ http://www.stsci.edu/resources/software_hardware/stsdas/synphot

Table 1. Estimation of the Zodiacal Light and Extragalactic Background Light for the BFM02 Field at $0.8\mu\text{m}$ using the ground-based ZL observations of Bernstein, Freedman & Madore (2002b) (BFM02), with corrections according to Mattila (2003) (M03). Systematic errors are given in brackets.

Quantity	Unit	Value	Comment
1 I_{tot}^λ (HST), 8000 Å	10^{-9} erg/s cm^2sr Å	$72.4 \pm 0.14, [\pm 1.0]$	BFM02, with “1- σ ” systematic error
2 I_{ZL}^λ ($0.80\mu\text{m}$)	10^{-9} erg/s cm^2sr Å	$69.4 \pm 0.4, [-0.9, +0.8]$	BFM02, with “1- σ ” systematic error
3 I_{ZL}^λ ($0.80\mu\text{m}$)	10^{-9} erg/s cm^2sr Å	$69.4 \pm 0.4, [-3.0, +1.6]$	BFM02, but with conservative systematic error
4 I_{ZL}^λ ($0.80\mu\text{m}$)	10^{-9} erg/s cm^2sr Å	56.3-60.0	BFM02 as corrected by M03
5 $I_{EBL+DGL}^\lambda$ ($0.80\mu\text{m}$)	10^{-9} erg/s cm^2sr Å	$3.0 \pm 0.4, [\pm 0.9]$	BFM02, with 1- σ statistical and “1- σ ” systematic errors
6 $I_{EBL+DGL}^\lambda$ ($0.80\mu\text{m}$)	10^{-9} erg/s cm^2sr Å	$3.0 \pm 0.8, [-1.9, +3.2]$	BFM02, but with 2- σ statistical and conservative systematic errors
7 $I_{EBL+DGL}^\lambda$ ($0.80\mu\text{m}$)	10^{-9} erg/s cm^2sr Å	$< (12.4 - 16.1)$	BFM02 with corrected ZL from M03
8 $\lambda \cdot I_{EBL+DGL}^\lambda$ ($0.80\mu\text{m}$)	nW/m ² sr	$< (99 - 129)$	BFM02 with corrected ZL from M03

Table 2. Estimation of the Zodiacal Light and Extragalactic Background Light for the BFM02 Field at $0.8\mu\text{m}$ using the Kelsall et al. (1998) (K98) Zodiacal Light model. Systematic errors are given in brackets.

Quantity	Unit	Value	Comment
1 I_{tot}^λ (HST), 8000 Å	10^{-9} erg/s cm^2sr Å	$72.4 \pm 0.14, [\pm 1.0]$	BFM02, with “1- σ ” systematic error
2 I_{ZL}^ν (model), J band	MJy sr^{-1}	$0.151, [\pm 0.0062]$	K98 model value from DSZA
3 Colour correction factor K		0.985	Assuming 6000 K blackbody
4 I_{ZL}^ν ($1.25\mu\text{m}$)	MJy sr^{-1}	$0.153, [\pm 0.0063]$	Monochromatic intensity
5 I_{ZL}^λ ($1.25\mu\text{m}$)	10^{-9} erg/s cm^2sr Å	$29.4, [\pm 1.15]$	Monochromatic intensity
6 $F_\odot^\lambda$ ($1.25\mu\text{m}$)	erg/s cm^2 Å	44.6	Monochromatic Solar flux
7 $F_\odot^\lambda$ ($0.80\mu\text{m}$)	erg/s cm^2 Å	112.3	Solar flux through WFPC f814
8 $F_\odot^\lambda$ ($0.80\mu\text{m}$)/ $F_\odot^\lambda$ ($1.25\mu\text{m}$)		2.52	Solar SED
9 I_{ZL}^λ ($0.80\mu\text{m}$)/ I_{ZL}^λ ($1.25\mu\text{m}$)		$2.26[\pm 0.05]$	Reddened Solar SED
10 I_{ZL}^λ ($0.80\mu\text{m}$)	10^{-9} erg/s cm^2sr Å	$66.4[\pm 2.6], [\pm 1.5]$	ZL resulting from the K98 model
11 $I_{EBL+DGL}^\lambda$ ($0.80\mu\text{m}$)	10^{-9} erg/s cm^2sr Å	$6.0 \pm 0.1, [\pm 3.2]$	I_{tot}^λ (HST) minus I_{ZL} (K98)
12 $\lambda \cdot I_{EBL+DGL}^\lambda$ ($0.80\mu\text{m}$)	nW/m ² sr	48.0 ± 25.6	$\lambda \cdot I_{tot}^\lambda$ (HST) minus $\lambda \cdot I_{ZL}$ (K98)

account for the ZL reddening relative to the Solar SED by applying a correction factor according to Eq.(22) of Leinert et al. (1998). The reddening correction $1.25\mu\text{m} \rightarrow 0.80\mu\text{m}$ of 1.116 is estimated to have a systematic error of ± 0.023 , i.e. $\pm 20\%$. The resulting ZL brightness of $66.4[\pm 2.6], [\pm 1.5]$ at $0.80\mu\text{m}$ is given in line 10 of Table 2. The first error in brackets corresponds to the Kelsall et al. (1998) model uncertainties, the second one to the 20% error adopted for the ZL reddening correction.

2.3 The residual signal, EBL + DGL, after subtraction of the ZL

BFM02a give for their *HST* measurement of I_{tot} at $0.80\mu\text{m}$ the value $72.2 \pm 0.14 [\pm 1.0] 10^{-9}$ erg/s cm^2sr Å (line 1 of Table 1 and Table 2). Their method of treating systematic errors leads to the paradoxical result (line 5 of Table 1) that subtracting the value of $I_{ZL}(0.8\mu\text{m}) = 69.4 \pm 0.4[-0.9, +0.8] 10^{-9}$ erg/s cm^2sr Å leaves as difference the value $I_{EBL+DGL}(0.8\mu\text{m}) = 3.0 \pm 0.4[\pm 0.9] 10^{-9}$ erg/s cm^2sr Å, i.e. the systematic error has *decreased* from its level for $I_{tot}(0.8\mu\text{m})$ alone. We give in line 6 of Table 1 a more conservative error estimate with 2σ statistical and quadratically added systematic errors. The systematic errors of $\sim \pm 5$

10^{-9} erg/s cm^2sr Å caused by the four corrections discussed by Mattila (2003) and in Sect. 2.1 above have not been included here. Instead, lines 7 and 8 give the estimated upper limits of $I_{EBL+DGL}^\lambda$ and $\lambda \cdot I_{EBL+DGL}^\lambda$, reflecting the conclusion of Mattila (2003) that the claim for a detection of the EBL by BFM02 was premature.

Subtraction of the Kelsall et al. (1998) I_{ZL} estimate from I_{tot} results in the values of $I_{EBL+DGL}^\lambda$ and $\lambda \cdot I_{EBL+DGL}^\lambda$ as given in lines 11 and 12 of Table 2. The systematic errors of I_{tot} and I_{ZL} have been added in quadrature.

Fig. 1 shows the $\lambda \cdot I_{EBL+DGL}^\lambda$ values and upper limits. They include the *IRTS* Near-Infrared Spectrometer (NIRS) values of Matsumoto et al. (2005) between $1.4 - 4\mu\text{m}$ and the *COBE*/DIRBE values at $1.25, 2.2, 3.5$, and $4.9\mu\text{m}$ as compiled in Dwek, Arendt, & Krennrich (2005) and Hauser & Dwek (2001). Our two estimates at $0.8\mu\text{m}$ are shown as an upper limit (box), based on the BFM02b ZL estimate as revised by Mattila (2003), and as an open circle with error bars, based on the ZL estimate from the Kelsall et al. (1998) model.

Table 3. Estimation of the diffuse Galactic surface brightness due to scattered starlight, $I_{DGL}(sca)$, and non-equilibrium emission or fluorescence by dust, $I_{DGL}(em)$.

Field	l, b deg	$N(H)$ 10^{20} cm^{-2}	A_V mag	λ μm	λI_{ISRF}^λ $\text{nW/m}^2\text{sr}$	A_λ mag	$\lambda I_{ISRF}^\lambda A_\lambda$	$\lambda I_{DGL}^\lambda(sca)$	$\lambda I_{DGL}^\lambda(em)$ $\text{nW/m}^2\text{sr}$	$\lambda I_{DGL}^\lambda(sca+em)$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
BFM02	206.6, -59.8	1.5	0.080	0.55	660 ^a	0.080	53	21 ^b	–	21
				0.80	867 ^a (900)	0.048	42	17 (8.0 ^c)	–	17
Matsumoto et al. (2005)	80-90, 42-48	1.3	0.069	1.25	1060 ^d	0.019	20	7.9	0.3	8.2
				1.65	1110 ^d	0.012	13	5.2	1.2	6.4
				2.2	655 ^d	0.0075	4.9	1.9	3.8	5.7
				3.5						5.0
				4.9						3.6

^a Mattila 1980^b derived from $I_{DGL}(sca)/I(100\mu\text{m})$ in L 1642 (Laureijs, Mattila & Schnur 1987)^c effective value to be applied to $I_{EBL+DGL}$ of BFM02^d Lehtinen & Mattila (1996)

3 THE CONTRIBUTION BY THE DIFFUSE GALACTIC LIGHT

We have so far presented in Tables 1 and 2 and Fig. 1 the combined contribution of the EBL and the DGL. Since Matsumoto et al. (2005) do not make any statement on the DGL they appear to have included it into their nominal “EBL” values. For the *COBE*/DIRBE based values at 1.25, 2.2, 3.5, and 4.9 shown in Fig. 1 the DGL has been considered to different degrees. The following discussion in this section is concerned with the DGL contribution to the Matsumoto et al. (2005) 1.4 – 4 μm and the BFM 0.80 μm values.

BFM02a used a general model of the ISRF and dust distribution to estimate the DGL (scattered starlight) contribution in their target field at 0.80 μm : $I_{DGL} = 0.8 \cdot 10^{-9} \text{ erg/s cm}^2\text{sr } \text{\AA}$. Mattila (2003) revisited the model estimate and argued that the dust column density toward the target field had been underestimated by a factor of ~ 3 and estimated the DGL intensity at 0.8 μm to be $I_{DGL} = 2.3 \cdot 10^{-9} \text{ erg/s cm}^2\text{sr } \text{\AA}$ corresponding to 18 $\text{nW/m}^2\text{sr}$. However, after including a correction caused by the fact that the DGL has a Fraunhofer line spectrum resembling the ZL spectrum Mattila (2003) found that an *effective* I_{DGL} correction of $\sim 1 \cdot 10^{-9} \text{ erg/s cm}^2\text{sr } \text{\AA}$ or 8 $\text{nW/m}^2\text{sr}$ shall be subtracted from the observed $I_{EBL+DGL}$ value of BFM02.

At optical and NIR wavelengths up to 1.5 μm the diffuse Galactic surface brightness is primarily due to Galactic starlight scattering off interstellar dust grains. At longer wavelengths ($\sim 2 - 5 \mu\text{m}$), because of decreasing optical depth of dust ($A_\lambda \lesssim 0.15 A_V$), the scattered light contribution rapidly decreases. However, recent *COBE*/DIRBE and *Spitzer* observations have shown that there is at NIR wavelengths, $\lambda > 2 \mu\text{m}$, a substantial diffuse Galactic surface brightness component (Bernard et al. 1994; Arendt et al. 1998; Arendt & Dwek 2003; Flagey et al. 2006) associated with the widely distributed low density dust medium. Such a NIR continuum emission had been previously detected in reflection nebulae by Sellgren, Werner & Dinerstein (1983). It can be due e.g. to non-equilibrium emission by very small transiently heated grains or to fluorescence

emissions by PAHs or very small grains; for a discussion see Flagey et al. (2006).

We will estimate the diffuse Galactic emission at the different wavelengths and for the two regions of sky targeted in BFM02 and Matsumoto et al. (2005) observations. The line-of-sight dust column density can be estimated using the 21-cm hydrogen column density map of Kalberla et al. (2005). For this map the best state-of-the-art stray radiation corrections have been applied, important for the high latitude fields with weak HI emission. The HI column densities for the two fields are 1.5 and 1.3 10^{20} cm^{-2} as given in column(3) of Table 3. Using the standard $N(H)/E(B - V)$ ratio (Bohlin, Savage, & Drake 1978) and $R_V = 3.1$ they correspond to the line-of-sight extinctions of $A_V = 0.080$ and 0.069 mag, respectively. We assume that the scattering properties of the dust grains, i.e. albedo a and forward scattering parameter g , are approximately constant between 0.55 and 2.2 μm (for justification see Lehtinen & Mattila 1996). Then, the scattered light intensity at different wavelengths is, for an optically thin line of sight towards high galactic latitudes, in a first approximation proportional to the product $I(\text{ISRF}) \times A_\lambda$. We give in columns (6)-(8) of Table 3 our estimates for $I(\text{ISRF})$ and A_λ . The wavelength dependence of A_λ corresponds to $R_V = 3.1$ (Mathis 1990). For $I(\text{ISRF})$ we have adopted at 1.25 – 2.2 μm the mean sky brightness values as obtained from the *COBE*/DIRBE Zodi Subtracted Mission Averaged (ZSMA) maps; see Lehtinen & Mattila (1996).

The scattered light intensity, $I_{DGL}(sca)$, has been measured in the translucent high latitude ($b = -36.7$ deg) cloud L 1642 as a function of optical thickness for five wavelengths between $\lambda = 0.35 - 0.55 \mu\text{m}$ (Laureijs, Mattila & Schnur 1987; Mattila 1990). At small optical depths we obtain from these observations at 0.55 μm the relation $I_{DGL}(sca)/I(100\mu\text{m}) = 3.0 \cdot 10^{-9} \text{ erg/s cm}^2\text{sr } \text{\AA} / \text{MJy sr}^{-1}$ which for $I(100\mu\text{m})/N(\text{HI}) = 0.85 \text{ MJy sr}^{-1}/10^{20} \text{ cm}^{-2}$ (Boulanger et al. 1996) corresponds to $I_{DGL}(sca)/N(\text{HI}) = 2.55 \cdot 10^{-9} \text{ erg/s cm}^2\text{sr } \text{\AA} / 10^{20} \text{ cm}^{-2}$. Thus, for the BFM02 field we obtain at 0.55 μm the estimate $\lambda I_{DGL}^\lambda(sca) = 21 \text{ nW/m}^2\text{sr}$ as given in column (9) of Table 3. The $\lambda I_{DGL}^\lambda(sca)$




Figure 1. Observed values and upper limits for the Extragalactic Background Light in the red and near infrared $\lambda = 0.8\text{--}5 \mu\text{m}$. The two estimates at $0.8 \mu\text{m}$ derived in this paper are indicated by an open circle and error bars (ZL model of Kelsall et al. 1998) and a rectangular box (BFM02 ZL observations with corrections according to Mattila 2003). The open circles with error bars at $1.4\text{--}4 \mu\text{m}$ are the *IRTS* Near-Infrared Spectrometer (NIRS) values of Matsumoto et al. (2005). The *COBE/DIRBE* values at 1.25 , 2.2 , 3.5 , and $4.9 \mu\text{m}$ are indicated by big solid symbols: squares (Dwek & Arendt 1998; Arendt & Dwek 2003), triangles (Cambr  sy et al. 2001), octagons (Wright 2001), and encircled octagons (Gorjian, Wright & Charry 2000). The integrated light from known galaxies is shown by small solid symbols: hexagons (Madau & Pozzetti 2000), triangles (Totani et al. 2001) and squares (Fazio et al. 2004). The estimated lower and upper limit for the *total* light of galaxies is shown by bars at 0.61 , 0.81 , 1.25 , and $2.2 \mu\text{m}$ (Totani et al. 2001). Our estimate for the Diffuse Galactic Light, $I_{DGL}(\text{sca}+\text{em})$, is shown as a dashed line according to column 11 of Table 3

values for the other wavelengths in Table 3 are then derived by scaling this $\lambda I_{DGL}^{\lambda}(\text{sca})$ at $0.55 \mu\text{m}$ with the $\lambda I_{ISRF}^{\lambda} \times A_{\lambda}$ values given in column (8). The value thus obtained for the BFM02 field at $0.80 \mu\text{m}$ is $17 \text{ nW/m}^2\text{sr}$ which very closely corresponds to the model estimate of BFM02a as corrected by Mattila (2003). These optical DGL estimates are in general terms also confirmed by the surface photometry of L 1780 (Mattila 1979) which stretched over the larger wavelength range of $0.35\text{--}0.755 \mu\text{m}$ but lacked the smallest optical depth range important for the present purpose. It can be seen from Table 3 column (9) that $\lambda I_{DGL}^{\lambda}(\text{sca})$ drops rapidly at $\lambda > 1.5 \mu\text{m}$.

We use the results of Flagey et al. (2006) to estimate the $\lambda I_{DGL}^{\lambda}$ values at $1.25\text{--}4.9 \mu\text{m}$ for the Matsumoto et al. (2005) field. In their Fig. 9 they give the diffuse Galactic

brightness at $2\text{--}14 \mu\text{m}$ for $N(\text{H}) = 10^{21} \text{ cm}^{-2}$. From this figure we read the $I_{DGL}(\text{tot}) = I_{DGL}(\text{sca}) + I_{DGL}(\text{emission})$ values for 2.2 , 3.5 , and $4.9 \mu\text{m}$ and they are given in column (11) of table 3. In order to estimate $I_{DGL}(\text{emission})$ at 1.25 and $1.65 \mu\text{m}$ we use the Flagey et al. (2006) model for the NIR continuum which they represent with a grey body of colour temperature $1100 \pm 300 \text{ K}$. The values thus obtained for $\lambda I_{DGL}^{\lambda}(\text{emission})$ are given in column (10) of Table 3. At these two wavelengths $I_{DGL}(\text{sca})$ already dominates over $I_{DGL}(\text{emission})$. It can be seen from Table 3 that the total $\lambda I_{DGL}^{\lambda}$ in units of $\text{nW/m}^2\text{sr}$ varies by a factor of ~ 2 over the NIR wavelength range, $\lambda = 1.25\text{--}4.9 \mu\text{m}$. At $0.80 \mu\text{m}$ it is ~ 2 times as large as at $1.25 \mu\text{m}$. However, after including the correction caused by the similarity of the DGL and ZL Fraunhofer line spectra its *effective* value, to be subtracted from the BFM02 $\lambda I_{EBL+DGL}^{\lambda}$ value at $0.80 \mu\text{m}$, becomes $8 \text{ nW/m}^2\text{sr}$, i.e. very similar to the value at $1.25 \mu\text{m}$.

The DGL contributions are, in general, substantially smaller than the $\lambda I_{EBL+DGL}^{\lambda}$ values of BFM02 and Matsumoto et al. (2005) shown in Fig. 1. The values are, however, closely similar to the integrated intensities derived from galaxy counts. Therefore, if the true NIR EBL surface brightness turns out to be of this order of magnitude the DGL contribution cannot be neglected. Rather, it will be the ultimate obstacle, even for space borne measurements from outside the Zodiacal cloud, which must be accurately known and subtracted before arriving at a credible measurement of the EBL.

4 DISCUSSION

4.1 Is there an $\sim 1 \mu\text{m}$ step in the EBL spectrum?

The interpretation of the NIR EBL excess as signature of Population III stars hinges on the reality of the claimed large discontinuity between 0.8 and $1.2\text{--}1.4 \mu\text{m}$. The values (upper limits) derived in Section 2 and shown in Fig. 1 for $\lambda I_{EBL+DGL}^{\lambda}$ at $0.8 \mu\text{m}$ are significantly larger than the original BFM02a value of $3.0 \cdot 10^{-9} \text{ erg/s cm}^2 \text{ sr } \text{\AA}$ or $24 \text{ nW/m}^2\text{sr}$ on which this conjecture was based.

The large range of the estimates at $0.80 \mu\text{m}$ reflects directly the differences and uncertainties of the values of $I_{ZL}(0.80 \mu\text{m})$ as derived from the Kelsall et al. (1998) model on the one, and from the BFM02b ground based observations on the other side. In both cases the ZL value has been extrapolated from another wavelength, $1.25 \mu\text{m}$ in the former and $0.465 \mu\text{m}$ in the latter case, using a reddened Solar SED to transform from Solar to ZL colour.

When assessing the reality of the $\sim 1 \mu\text{m}$ step in the EBL spectrum it is obvious that the subtraction of the foreground ZL has to be done using consistent ZL values, both on the short and on the long wavelength side of the step. With this in mind we have applied the Kelsall et al. (1998) model and it can be seen from Fig. 1 that there is, within the error limits, no evidence for the step. Errors in the basic assumptions or parameter values of the Kelsall et al. (1998) model would influence the values at $0.80 \mu\text{m}$ and at $> 1 \mu\text{m}$ in a similar way and are not expected to artificially create or destroy a large spectral step, if existent, between these wavelengths.

When applying the BFM02b measurement of I_{ZL} an extrapolation of their measured ZL value from $0.465 \mu\text{m}$ to

0.80 μm is needed. This introduces an uncertainty of similar size as the extrapolation of the Kelsall et al. (1998) model from 0.80 to 1.25 μm . Given the corrections needed for the BFM02b ZL measurement and adopting conservative systematic error estimates as discussed in Sect. 2.1 above, there is no evidence for a downward step in the EBL spectrum from 1.25 to 0.80 μm .

The Diffuse Galactic Light correction as discussed in Sect. 3 is of similar strength both below and above 1 μm and it does not, therefore, contribute to the possible discontinuity at $\sim 1 \mu\text{m}$.

The author wishes to emphasise that he does not advocate the (very) large $I_{\text{EBL}+\text{DGL}}$ value at 0.80 μm , apparently resulting from the discussion in this paper, as anything else but an upper limit. Rather, also the value derived using the Kelsall et al. (1998) ZL model, which appears to be different from zero, should be considered only as an upper limit.

4.2 Remarks on the near infrared EBL measurements

The analysis presented in this paper does not allow conclusions concerning the *overall level* of the NIR EBL at 1 - 4 μm . Given the systematic uncertainties of the ZL subtraction a direct photometric detection of the NIR EBL appears very difficult.

The Kelsall et al. (1998) model appears to give a good representation of the ZL thermal emission at mid and far IR wavelengths where the ZL dominates the sky brightness. However, at the near IR wavelengths, $1 \mu\text{m} < \lambda < 5 \mu\text{m}$, the Zodiacal component is weaker and its modelling relies on the mid IR results. As stated by Kelsall et al. (1998) their ZL model is not unique but the predicted values depend on the assumed models for the IPD cloud geometry and dust parameters. Furthermore, the NIR ZL is caused by scattering instead of thermal emission which means that dust parameters different from those at the mid IR wavelengths have to be used.

An alternative ZL model has been presented by Wright (1998) and has been applied in the Wright & Reese (2000), Wright (2001), and Gorjian, Wright & Charry (2000) analyses of the DIRBE data. The resulting EBL values are by ~ 33 , 8, and 4 $\text{nW/m}^2\text{sr}$ smaller at the J, K, and L bands, respectively, as compared to the results based on the Kelsall et al. (1998) model. This does not, however, substantially change the dilemma with an unexpectedly large NIR EBL value.

Dwek, Arendt, & Krennrich (2005) have presented arguments suggesting that the Matsumoto et al. (2005) EBL excess emission at $\lambda = 1.4 - 4 \mu\text{m}$ may be due to insufficient ZL subtraction. This possibility is especially suggested by the practically identical SEDs of the ZL and the “EBL excess” over the whole wavelength range of the observations, see Fig. 6 of Dwek, Arendt, & Krennrich (2005).

The “EBL excess” emission as derived from the *IRTS* data by Matsumoto et al. (2005) contains the contribution by the Diffuse Galactic emission. The DGL has been considered in several of the analyses of the *COBE*/DIRBE data (see e.g. Arendt et al. 1998; Arendt & Dwek 2003) but its separation from the much larger component due to the Integrated Starlight has been problematic because of the large field of view of DIRBE. The recent *Spitzer* observations

(Flagey et al. 2006) have offered a better estimate for the 2-5 μm DGL and they suggest that it is substantially larger than the values previously found by Arendt et al. (1998). Although the DGL level remains insignificant over the wavelength range 1.25-2.5 μm this is no longer the case at 3.5 μm . At this wavelength the sum of the integrated light from resolved galaxies (Fazio et al. 2004) and the DGL (Table 3) amounts to $\sim 10.4 \text{ nW/m}^2\text{sr}$ which is within its estimated error limits of $\sim \pm 2 \text{ nW/m}^2\text{sr}$ fully compatible with the Matsumoto et al. (2005) EBL value of $14.4 \pm 3 \text{ nW/m}^2\text{sr}$. In addition, similarly as at the shorter wavelengths (Totani et al. 2001), some contribution from the unresolved galaxies should be added (Kashlinsky et al. 2005; Savage & Oliver 2005) bringing the two estimates even closer to each other.

Recently, an upper limit to the EBL flux around 1 - 2 μm has been announced by Aharonian et al. (2006) using *HESS* observations of intergalactic absorption of γ -ray emission of blazars: $I_{\text{EBL}} \leq (14 \pm 4) \text{ nW/m}^2\text{sr}$. Thus, these *HESS* observations would suggest that more than two thirds of the EBL in the NIR band is resolved into individually detected galaxies. The *HESS* result depends, however, on the adopted TeV spectra of the blazars. Therefore, direct photometric observations of the optical/NIR EBL are still urgently needed.

5 CONCLUSIONS

We have scrutinised the observational evidence for the claimed discontinuity at $\sim 1 \mu\text{m}$ in the spectrum of the Extragalactic Background Light. We have also estimated the contribution of the Diffuse Galactic emission to the “EBL excess”. Our conclusions are the following:

- (1) The reality of the $\sim 1 \mu\text{m}$ step hinges on the claimed detection of the EBL by BFM02 at 0.80 μm . We find that after applying corrections to the analysis of their Zodiacal Light measurement only an upper limit to the EBL can be set which does not imply any spectral step between 0.80 and 1.25 μm .
- (2) An alternative estimate of the Zodiacal Light at 0.80 μm is obtained using the Kelsall et al. (1998) model. Again, the upper limit set to the EBL at 0.80 μm does not warrant any spectral step between 0.8 and 1.25 μm .
- (3) An estimate for the contribution by Diffuse Galactic emission is obtained from recent *Spitzer* measurements by Flagey et al. (2006). It is found that the sum of the Diffuse Galactic emission and the integrated light of resolved galaxies can explain the claimed EBL signal in the 3.5 μm window where the ZL contamination is at minimum.

REFERENCES

- Aharonian F. et al., 2006, *Natur*, 440, 1018
- Arendt, R.G. et al., 1998 *ApJ*, 508, 74
- Arendt R. G., Dwek E., 2003, *ApJ*, 585, 305
- Bernard, J.P. et al., Boulanger F., Desert F. X., Giard M., Helou G., Puget J. L., 1994, *A&A*, 291, L5
- Bernstein, R.A., Freedman, W.L., Madore, B.F., 2002a, *ApJ*, 571, 56
- Bernstein, R.A., Freedman, W.L., Madore, B.F., 2002b, *ApJ*, 571, 85

- Bernstein, R.A., Freedman, W.L., Madore, B.F., 2005, ApJ, 632, 713
- Bohlin R. C., Savage B. D., Drake J. F., 1978, ApJ, 224, 132
- Boulanger F., Abergel A., Bernard J.-P., Burton W. B., Desert F.-X., Hartmann D., Lagache G., Puget J.-L., 1996, A&A, 312, 256
- Cambr  sy, L., Reach, W.T., Beichman, C.A., Jarret. T.H., 2001, ApJ, 555, 563
- Colina, L., Bohlin, R.C., Castelli, F. 1996, AJ, 112, 307
- Cooray A., Yoshida N., 2004, MNRAS, 351, L71
- Dwek E., Arendt R. G., 1998, ApJ, 508, L9
- Arendt R. G., Dwek E., 2003, ApJ, 585, 305
- Dwek E., Arendt R. G., Krennrich F., 2005, ApJ, 635, 784
- Fazio G. G., et al., 2004, ApJS, 154, 39
- Flagey N., Boulanger F., Verstraete L., Miville Desch  nes M. A., Noriega Crespo A., Reach W. T., 2006, A&A, 453, 969
- Gorjian, V., Wright, E.L., Charry, R.R., 2000, ApJ, 536, 550
- Hauser M. G., Dwek E., 2001, ARA&A, 39, 249
- Kalberla, P. M. W.; Burton, W. B.; Hartmann, Dap; Arnal, E. M.; Bajaja, E.; Morras, R.; P  ppel, W. G. L., 2005, A&A 440, 775
- Kashlinsky A., Arendt R. G., Mather J., Moseley S. H., 2005, Natur, 438, 45
- Kelsall, T. et al. 1998 ApJ, 508, 44
- Laureijs R. J., Mattila K., Schnur G., 1987, A&A, 184, 269
- Lehtinen K., Mattila K., 1996, A&A, 309, 570
- Leinert, C. et al., 1998, A&AS 127, 1
- Longair M., 2001, IAUS, 204, 505
- Madau, P., Silk, J., 2005, MNRAS, 359, 37
- Madau P., Pozzetti L., 2000, MNRAS, 312, L9
- Mathis J. S., 1990, ARA&A, 28, 37
- Mattila K., 1979, A&A, 78, 253
- Mattila K., 1980, A&AS, 39, 53
- Mattila K., 1990, IAUS, 139, 257
- Mattila, K., 2003, ApJ, 591, 119
- Matsumoto, T. et al., 2005 ApJ, 626, 31
- Salvaterra R., Ferrara A., 2003, MNRAS, 339, 973
- Santos M. R., Bromm V., Kamionkowski M., 2002, MNRAS, 336, 1082
- Savage, R. S., Oliver, S., 2005 (astro-ph/0511359)
- Sellgren, K., Werner, M. W., Dinerstein, H. L., 1983, ApJ, 271, L13
- Totani T., Yoshii Y., Iwamuro F., Maihara T., Motohara K., 2001, ApJ, 550, L137
- Wright E. L., 1998, ApJ, 496, 1
- Wright, E.L., 2001, ApJ, 553, 538
- Wright E. L., Reese E. D., 2000, ApJ, 545, 43